

**PEMANFAATAN LIMBAH CAIR TEMPE DALAM MEDIA AIR
BUDIDAYA UNTUK PERTUMBUHAN DAN EFISIENSI IKAN NILA
(*Oreochromis niloticus*)**

NASKAH PUBLIKASI



Oleh :

HERYANTO
201610260311031

**FAKULTAS PERTANIAN PETERNAKAN
UNIVERSITAS MUHAMMADIYAH MALANG**

2020

**PEMANFAATAN LIMBAH CAIR TEMPE DALAM MEDIA AIR
BUDIDAYA UNTUK PERTUMBUHAN DAN EFISIENSI IKAN NILA**

(Oreochromis niloticus)

NASKAH PUBLIKASI

**Diajukan Kepada Universitas Muhammadiyah Malang Sebagai Salah Satu
Persyaratan Untuk Memperoleh Gelar Sarjana Perikanan**



Disusun Oleh :

HERYANTO

201610260311031

**FAKULTAS PERTANIAN PETERNAKAN
UNIVERSITAS MUHAMMADIYAH MALANG**

2020

HALAMAN PERSETUJUAN

PEMANFAATAN LIMBAH CAIR TEMPE DALAM MEDIA AIR
BUDIDAYA UNTUK PERTUMBUHAN DAN EFISIENSI IKAN NILA
(*Oreochromis niloticus*)

Oleh:

Heryanto
NIM : 201610260311031

Disetujui oleh :

Pembimbing Utama

Tanggal,



Anis Zubaidah, S.Pi., M.Si
NIDN. 0727028605

Pembimbing Pendamping

Tanggal,



Soni Andriawan, S.Pi., MP
NIDN. 0712069202

Malang.....

Menyetujui :

An. Dekan,
Wakil Dekan 1,



Dr. Ir. Aris Winaya, MM, M.Si
NIP. 196405.14199.003.1.002



Wakil Ketua Jurusan

Sanjaya Andriwirawan S., S.Pi, MP
NIP. 1971031410.0538

SKRIPSI

**PEMANFAATAN LIMBAH CAIR TEMPE DALAM MEDIA AIR
BUDIDAYA UNTUK PERTUMBUHAN DAN EFISIENSI IKAN NILA**

Oleh:


Heryanto

NIM : 201610260311031

Disusun berdasarkan Surat Keputusan Dekan Fakultas Pertanian Peternakan
Universitas Muhammadiyah Malang Nomor :
dan rekomendasi Komisi Skripsi Fakultas Pertanian Peternakan UMM pada
tanggal : dan keputusan Ujian sidang yang
dilaksanakan pada tanggal : 18 Agustus 2020

Dewan Penguji :

Pembimbing Utama




Anis Zubaidah, S.Pi., M.Si
NIDN. 0727028605

Pembimbing Pendamping



Soni Andriawan, S.Pi., MP
NIDN. 071501701

Penguji Utama



Ganjar Adhivirawan S., S.Pi., MP
NIP. 110.1410.0538

Penguji Kedua



Dr. Ir. David Hermawan, M.P., IPM
NIP. 196405261990031003

Malang.....

Mengesahkan :



Dekan,



Dr. Ir. David Hermawan, M.P., IPM
NIP. 196405261990031003



Ketua Jurusan,



Ganjar Adhivirawan S., S.Pi., MP
NIP. 110.1410.0538

SURAT PERNYATAAN

Yang bertanda tangan di bawah ini :

Nama : Heryanto
NIM : 201610260311031
Fakultas/Jurusan : Pertanian Peternakan/Perikanan
Perguruan Tinggi : Universitas Muhammadiyah Malang

Menyatakan dengan sebenarnya dan sesungguhnya, bahwa skripsi atau karya ilmiah yang berjudul :

Pemanfaatan Limbah Cair Tempe dalam Media Air Budidaya untuk Pertumbuhan dan Efisiensi Pakan Ikan Nila (*Oreochromis niloticus*)

1. Skripsi ini adalah milik saya sendiri yang disusun berdasarkan serangkaian penelitian yang saya lakukan dan belum pernah diajukan untuk memperoleh gelar pada program sejenis sejenis di perguruan tinggi manapun, semua data dan informasi yang digunakan telah dinyatakan secara jelas dan dapat diperiksa kebenarannya.
2. Penulisan skripsi ini tidak ada plagiasi, duplikasi ataupun replikasi terhadap hasil penelitian ini dari pihak-pihak manapun yang menyebabkan hasil penelitian ini tidak otentik, kecuali secara tertulis diacu dalam skripsi dan disebutkan rujukannya dalam daftar pustaka.
3. Skripsi ini disusun berdasarkan persetujuan dan bimbingan dari dewan pembimbing dan telah diuji dihadapan dewan penguji tugas akhir Jurusan Perikanan Fakultas Pertanian Peternakan Universitas Muhammadiyah Malang.

Demikian surat pernyataan ini saya buat dengan sebenar-benarnya dan bertanggung jawab terhadap pernyataan ini.

Malang, 18 September 2020

Mengetahui

Dosen Pembimbing Utama

Anis Zubaidah S.Pi., M.Si
NIDN. 0727028605

Yang menyatakan

Heryanto
NIM. 201610260311031

KATA PENGANTAR

Puji syukur penulis panjatkan kehadiran Allah SWT yang telah melimpahkan rahmat dan hidayahNya sehingga penulis dapat menyelesaikan skripsi dengan judul **Pemanfaatan Limbah Cair Tempe dalam Media Air Budidaya untuk Pertumbuhan dan Efisiensi Pakan Ikan Nila (*Oreochromis niloticus*)** sebagai salah satu syarat untuk memperoleh gelar sarjana perikanan di Universitas Muhammadiyah Malang.

Dalam proses penyusunan skripsi ini, penulis banyak mendapatkan bimbingan dan petunjuk serta bantuan yang bermanfaat dari berbagai pihak. Oleh karena itu, dalam kesempatan ini penulis ingin menyampaikan ucapan terima kasih yang sebesar-besarnya kepada :

1. Bapak Dr. Ir. David Hermawan M.P., IPM selaku dekan Fakultas Pertanian Peternakan Universitas Muhammadiyah Malang.
2. Ibu Anis Zubaidah S.Pi., M.Si dan Soni Andriawan, S.Pi., M.Sc selaku Pembimbing I dan Pembimbing II yang telah banyak meluangkan waktu dan pikiran untuk memberikan bimbingan dan arahan yang sangat berguna, hingga penulis dapat menyelesaikan perkuliahan dan skripsi ini dengan baik.
3. Bapak Ganjar Adhywirawan Sutarjo S.Pi., M.Si selaku ketua program studi Budidaya Perairan Universitas Muhammadiyah Malang sekaligus dosen wali yang telah mendukung dan memberi pengarahan sejak awal perkuliahan hingga selesainya skripsi ini.
4. Ayah dan Ibu, yang selalu selalu mendoakan penulis serta curahan kasih sayang yang tiada tara. Hal ini merupakan kekuatan terbesar bagi penulis untuk terus memiliki motivasi dalam perkuliahan dan proses skripsi ini.
5. Teman-teman Jurusan Perikanan khususnya angkatan 2016 kelas A yang selalu memberikan semangat dan juga membantu proses penelitian penulis
6. Laboratorium Perikanan beserta rekan-rekan asisten, untuk setiap dukungan dan bantuan selama ini.
7. Semua pihak yang tidak dapat penulis sebutkan satu persatu, yang telah banyak memberikan bantuan pada penulis dalam menyelesaikan skripsi ini

Penulis menyadari tiada satupun karya manusia yang sempurna, sehingga kritik dan saran demi perbaikan karya ini sangat penulis harapkan. Meski demikian, penulis berharap semoga ini dapat bermanfaat bagi peneliti khususnya dan pembaca pada umumnya.

Malang, 18 September 2020

Penulis

Heryanto

DAFTAR ISI

COVER	i
HALAMAN PERSETUJUAN	iii
HALAMAN PENGESAHAN	iv
SURAT PERNYATAAN	v
KATA PENGANTAR	vi
DAFTAR ISI	vii
DAFTAR TABEL	viii
DAFTAR GAMBAR	ix
DAFTAR LAMPIRAN	x
ABSTRAK	1
PENDAHULUAN	1
METODE PENELITIAN	2
Waktu dan Tempat	2
Alat dan Bahan	2
Metode	3
Prosedur	3
Persiapan Wadah	3
Hewan Uji	3
Persiapan Media	3
Parameter Uji	4
Pertumbuhan Mutlak	4
Laju pertumbuhan Spesifik Harian	4
Pertumbuhan Panjang Mutlak	5
Pemanfaatan Pakan	5
Kualitas Air	5
HASIL DAN PEMBAHASAN	5
Pertumbuhan Berat Mutlak, Panjang Mutlak, Pertumbuhan Harian	5
Efisiensi Pemanfaatan Pakan	7
Kualitas Air	8
KESIMPULAN DAN SARAN	9
UCAPAN TERIMA KASIH	9
DAFTAR PUSTAKA	10
LAMPIRAN	

DAFTAR TABEL

Tabel	Halaman
1. Hasil Berat Mutlak, Berat spesifik harian, Panjang Mutlak	5
2. Hasil pengamatan kualitas air	8



DAFTAR GAMBAR

Gambar	Halaman
1. Grafik Efisiensi Pemanfaatan Pakan	8



DAFTAR LAMPIRAN

Gambar

Halaman

Form Hasil Cek Plagiasi Tugas Akhir 12



Utilization of Tempeh Liquid Waste in Cultural Water Media for Growth and Feed Utilization Efficiency on Nile tilapia (*Oreochromis niloticus*)

Heryanto^{1a}, Anis Zubaidah^{1,b}, Soni Andriawan^{1,c,*}.

Department of Fisheries, Faculty of Agriculture and Animal Science,
University of Muhammadiyah Malang, Malang East Java 65144, Indonesia.

Correspondence Email: soniandriawan1992@gmail.com

ABSTRACT. Tempeh waste has many nutrients including protein, fat, carbohydrates, Nitrogen Phosphorus, potassium and also has positive microbes which, if processed, can be used as utilization in fish farming. The purpose of this study is to determine whether the liquid waste media in tempeh waste can be utilized in the growth of tilapia (*Oreochromis niloticus*) and can make efficient use of feed. Test animals used were as many as 120 tilapia fish 5-8 cm. Tempeh waste used is liquid tempe waste that is the result of boiling soybeans. This research method uses a Completely Randomized Design (CRD) with treatment P0 (without liquid tempeh waste) P1 (5% waste dose) P2 (10% waste dose) and P3 (15% waste dose) by dissolving the dose into the fish culture media. 1.42 ± 0.22 , GR (8.61 ± 1.45) and absolute length (1.82 ± 0.38) . The best EPP is P2 treatment (12.92 ± 6.77) and the lowest is P0 (4.77 ± 3.05) . Tempeh liquid waste in the media of tilapia (*Oreochromis niloticus*) significantly influences ($P < 0.05$) on absolute growth but does not significantly affect the specific daily weight gain and absolute length as well as on feed efficiency.

Keywords: Tempeh Liquid Waste, Growth, Efficiency of Feed Utilization, Ammonia.

INTRODUCTION. Tempeh industry is a business that is able to influence the economic movement of the community. The existence of the tempe industry is not only in the environment of big cities but also in small cities. In Indonesia, there are currently around 81 thousand tempe industries that produce 2.4 million tons of tempe per year (BSN, 2012). The presence of tempeh industry in addition to having a positive impact also has a negative impact, namely the presence of waste that is not easy to process. The process of producing tempe produces solid and liquid waste. Liquid waste in processed tempe comes from soaking water, boiling, washing and stripping. The handling of wastewater produced cannot currently be utilized by the community and the resulting liquid waste is discharged into sewers and rivers, in contrast to solid waste, solid waste can be used as animal feed such as cattle and goats. Liquid waste discharged in the river can be polluted this because the nature of the liquid waste produced from tempe has an odor other than that algae blooms can occur in the waters due to organic compounds.

The compounds contained in tempe consist of 0.42% protein, fat 0.13%, carbohydrate 0.11%, water 98.87%, calcium 13.60 ppm, phosphorus 1.74 ppm and iron 4.55 ppm. If used appropriately, it will reduce environmental pollution

and eliminate the source of disease (Said, 1999). The contents in tempe liquid waste include nitrogen gas (N²), oxygen (O²), hydrogen sulfide (H²S), Ammonia (NH³), carbon dioxide (CO²), and methane (CH⁴) (Said 1999). In addition to the content of organic matter contained in tempe waste there are also bacteria. In the research "Isolation and Characterization of Lactic Acid Bacteria from Soybean Soaking Liquid Waste" obtained eight isolates that were successfully isolated from soybean soaking liquid waste, seven of which were identified as Lactic Acid Bacteria (LAB) which were suspected as members of the genus *Lactobacillus* (Amaliah *et al.* 2018) .

Lactic Acid Bacteria (LAB) is a type of gram-positive bacteria that has a role as a probiotic there are various beneficial microorganisms, namely *Lactobacillus* which is useful for fermenting organic material into lactic acid compounds, photosynthetic bacteria that function to absorb toxic gases and heat from the fermentation process, yeast which has a role in fermenting organic material into alcohol compounds, sugar and amino acids and *Actinomycetes* that function to produce antibiotic compounds that are toxic to pathogenic bacteria and able to dissolve phosphate ions and other micro ions (Wididana, 1993).

Tilapia is a fish that currently has a high economic value, and is a freshwater fish that is relatively easy to cultivate because it has a tolerant of the water is quite high. Besides tilapia has a high protein so that in addition to a good taste the nutritional value of tilapia will also provide a reason for people to consume tilapia, so high consumption of tilapia will increase market demand until now the development of tilapia production from year on year has increased in 2017 tilapia fish production reached 1.15 million tons, up by 3.6 percent from 2016 which reached 1.14 million tons (KKP 2017).

RESEARCH METHODS

Time and Place

This research was conducted on January 16 to February 16, 2020 which took place at the Fisheries Laboratory, Faculty of Agriculture-Animal Husbandry, Muhammadiyah University of Malang. East Java - Indonesia.

Material

The tools used in this study were 12 aquariums, analytical scales, rulers, pH meters, DO meters, thermometers, basins, nets, blowers, jerry cans, and aeration hoses. The materials used in this study are pellet feed, 5-6 cm tilapia fish, tempeh liquid waste and fresh water.

Methods

The method used in this study is a non factorial completely randomized design (RAL) experimental method. The type of factor tested was the difference in tempe liquid waste dosage consisting of four treatments and each consisting of 4 replications. The experimental unit is in the form of a 20x30x30 aquarium with 12 units with each aquarium filled with 25 liters of water and 10 fishes. The test treatments are as follows:

- Treatment P0: Tempeh liquid waste 0% per liter

- Treatment P1: Tempeh liquid waste 5% per liter
- Treatment P2: Tempeh liquid waste 10% per liter
- Treatment P3: Tempeh liquid waste 15% per liter

Procedures

Container Preparation

The container used in this study was an aquarium measuring 20x30x30 cm. The container used previously was washed first with water and sunlight then sterilized with a backlin to remove mold and bacteria. After that the aquarium is rinsed with clean water and then dried for 1 day, then each aquarium is filled with 25 liters of water and aerated.

Test Animals

Test animals used in this study were 5-8 cm tilapia obtained from the Pendem Fish Seed Center, Junrejo District, Batu City, East Java. The fish is put into a fiber bath for acclimation for 3 days.

Media preparation

The medium used is fresh water from the water pump of the fisheries lab and also as a test material, namely tempe liquid waste originating from the tempe industry in the village of Sanan, Malang. The tempe liquid waste is then mixed with the water of each aquarium and according to the concentration of the treatment. Then the media is allowed to stand for 3-7 days to reduce the level of wastewater saturation.

Test Parameters

Absolute Growth

Absolute growth or weight gain can be calculated using the formula (Dewantoro, 2001) as follows:

$$W = W_t - W_0$$

Information :

- W = Absolute weight growth (gr)
- W_t = Weight of the final fish rearing (gr)
- W₀ = Weight of the initial fish in the study (gr)

Specific Growth Rate Daily or Specific Growth Rate

According to Agustin et al, (2014), the formula for calculating specific growth rates is:

$$SGR = \frac{\ln W_t - \ln W_0}{t} \times 100\%$$

Information :

- SGR = Daily growth rate (%)
- W₀ = Average seed weight at the beginning of the study (gr)
- W_t = Average weight of seeds on the t-day (gr)
- t = Maintenance time (days)

Absolute Length Growth

Absolute length growth can be calculated using the formula (Effendi *et al.* 2006), namely:

$$L = L_t - L_0$$

Information :

L. = Growth in absolute length (cm)

L_t = Final length (cm)

L₀ = Initial length (cm)

Efficiency of Feed Utilization

The daily relative growth rate of fish is calculated using the NRC formula (1993) as follows:

$$EPP = \frac{W_t - W_0}{F} 100\%$$

Information :

EPP = Efficiency of feed utilization

F = Amount of feed (g)

W_t = Animal weight of the final test of the study (g)

W₀ = Weight of the initial research animal (g)

Water quality

The quality of water observed in this study includes the level of water acidity (pH), temperature, Dissolved oxygen (D_O) and ammonia. Data collection related to water quality is carried out routinely, once a week.

Data Collection Techniques

Sampling of data is done randomly (*Random sampling*).

To find out the growth of fish, 50 percent of the fish stocking population was tested for fish. Sampling of fish is done regularly, that is, once a week.

RESULTS AND DISCUSSION

Growth in absolute weights, Specific length and daily growth

The results of the analysis showed that the highest absolute weight growth was at a concentration of 5% per liter (P1) of 8.61 grams while the lowest absolute weight value was at 0% per liter (P0) of 3.42 grams. The absolute length increase of the research is highest in the P1 1.82 treatment and the lowest in the P0 1.22 treatment. The highest daily weight gain in tilapia during the study was in the treatment P1 of 1.5 g and the lowest in the treatment of P0 was 0.73. Data on absolute weight growth, daily growth rate and absolute length during maintenance can be presented in Table 1.

Table.1 Absolute weight (GR), Specific daily growth (SGR) and Absolute length.

Parameter	P0 (0% / liter)	P1 (5% / liter)	P2 (10% / liter)	P3 (15% / liter)
Absolute Weight (W)	3.42 ± 2.06	8.61 ± 1.45	5.70 ± 1.12	4.48 ± 0.68
SGR (% / day)	0.73 ± 0.45	1.5 ± 0.23	1.26 ± 0.26	1.21 ± 0.25
Absolute Length (cm)	1.22 ± 0.34	1.82 ± 0.38	1.54 ± 0.30	1.67 ± 0.34

The results of the analysis carried out using diversity (ANNAVA) F count $> F$ table 1% and 5% so that it can be said the absolute growth value is significantly different. The results of the LSD P0 test were significantly different from P1 and were not significantly different from P2 and P3. P1 is significantly different from P0, P2 and P3. P2 is not significantly different from P0 and P3 but significantly different from P1. P3 is not significantly different in P0 and P2 but significantly different in P1. However, for absolute length and growth of specific weights from the results of diversity (ANNAVA) F count $< F$ table 1% and 5% so that it can be said not significantly different. From the absolute growth value data table the highest value is in the treatment of tempe liquid waste 5% per liter (P1) and the lowest is P0 or no treatment. Likewise with the absolute length and the highest daily weight gain is the treatment P1 and the lowest is P0.

The difference in length and weight that is not the same has a significant effect on this because the growth of fish weight is faster than the increase in length, same with the opinion of Islaminingrum (2011) in Mashuri *et al.*, (2012) that the relationship between length and weight that occurs in fish, is that positive allometric which shows that the growth of fish weight is faster than the growth in length and negative allometric which shows that the growth of fish length is faster than the growth in weight. This difference in growth value is caused because liquid waste from tempe contains organic material such as N, P and K, besides that in tempe waste there are positive bacteria that are bacteria that can benefit organism. Nurosid, (2011) states that the largest component of tempe liquid waste is protein (N-total) of 226.06 mg / L to 434.78 mg / L, so that the entry of tempe liquid waste into the aquatic environment will increase the total nitrogen in these waters. MZ (2015) states that organic material that enters the waters will be decomposed into nutrients, for example nitrogen (N) and phosphorus (P). Andersen *et al.* (2006) stated that an increase in nutrients in water, especially N and P, can trigger the growth of phytoplankton and aquatic plants which can affect the structure, function and balance of the ecosystem. The concentration of these nutrients will be able to make the waters fertile, with fertile waters, the availability of nutrients for fish will be fulfilled so that growth will be good.


The types of bacteria found in tempe liquid waste are Lactic Acid Bacteria (LAB). In the research of Amaliah *et al* (2013) regarding the isolation and characterization of Lactic Acid Bacteria from Soybean Tempe Wastewater, eight isolates were isolated, seven of which were identified as lactic acid bacteria belonging to the genus *Lactobacillus*. Sulistijowati (2016) states that lactic acid bacteria are gram-positive and false-negative bacteria that produce lactic acid in carbohydrate fermentation, and are normal flora found in the digestive tract of animals, both land and aquatic animals. Wulandari (2017) said that lactic acid bacteria have an important role in fish life, because naturally these bacteria are able to boost the immune system and help the digestive organs in digestion. Lactic

acid bacteria tend to be able to increase specific and nonspecific immune systems, such as lysozyme activity in the body, macrophages, and respiratory bursts. Lactic Acid Bacteria enter the digestive tract of fish and live in it. Bacteria in digestion then secrete digestive enzymes such as protease and amylase enzymes (Irianto, 2003). Gatescoupe (1999) states that the enzyme can function as a catalyst in the digestion of carbohydrates and proteins. In addition, the function of LAB plays an important role in breaking down organic matter in a waters so that it can improve water quality in a media. Hadioetomo, (1993) a group of lactic acid bacteria that is in a digestion will act as a probiotic and if it is present in an aquatic environment it acts as a decomposer

Feed Utilization Efficiency

The results of the study of the addition of tempe liquid waste to the efficiency of tilapia fish utilization can be seen in figure 1.

Graphic Figure 1. Efficiency of Tilapia Feed Utilization during the study (%).



The results of the analysis carried out using diversity (ANNOVA) F count <F table 1% and 5% so that it can be said the efficiency of feed utilization is not significantly different. Even though the treatment was not significantly different, the lowest value of the utilization of feed was in the treatment P0 (0%) compared to other treatments the Feed Utilization Efficiency value was 4.77 ±. Feed Utilization Efficiency is the most important factor in supporting fish growth. Feed quality, amount of feed and water quality greatly affect the growth of fish weights. The low value of treatment P0 (0%) is due to the alleged lack of nutrients in the media and the absence of gram-positive microbial bacteria that can help the digestive process. Statement by Widanarni *et al.* (2009) that feed efficiency in the treatment with the application of biofloc technology is higher because of an increase in biofloc biomass as a source of nutrition or supplementary food for cultivation cultivation. The abundance of microbes in aquaculture media is also expected to play an active role in the digestive process such as the application of probiotics in fish feed. From the results of the study of Ahmadi *et al.* (2012) giving probiotics at a dose of 108 cells / ml only gave an efficient value of feed utilization of 43.93%. The high value of the efficiency of the use of feed given tempe liquid waste in tilapia culture media shows a better value compared to the

treatment without tempe wastewater treatment. The abundance of microbes in aquaculture media is also expected to play an active role in the digestive process such as the application of probiotics in fish feed. From the results of the study of Ahmadi *et al.* (2012) giving probiotics at a dose of 108 cells / mL only gave an efficient value of feed utilization of 43.93%. The high value of the efficiency of the use of feed given tempe liquid waste in tilapia culture media shows a better value compared to the treatment without tempe wastewater treatment. The abundance of microbes in aquaculture media is also expected to play an active role in the digestive process such as the application of probiotics in fish feed. From the results of the study of Ahmadi *et al.* (2012) giving probiotics at a dose of 108 cells / mL only gave an efficient value of feed utilization of 43.93%. The high value of the efficiency of the use of feed given tempe liquid waste in tilapia culture media shows a better value compared to the treatment without tempeh wastewater treatment.

Water quality

Data on water quality results during the study can be presented in table 2.

Table 2. Water Quality During the Study

No Treatment		Water Quality Parameters			
		Temperature	pH	DO (mg / l)	Ammonia The beginning
1	P0	25-29	6.5-8.8	2.0-2.9	0.0-1,5
2	P1	25-29	6,9-8,7	2.1-4.0	0.5-0.25
3	P2	25-29	7,1-8,6	2.2-3.3	1.5-0.25
4	P3	25-29	7,3-8,6	2.2-3.1	3.0-0.25

The degree of acidity obtained during the study of all treatments ranged from 6.5 to 8.8. From the pH data obtained it can be seen that the pH of all treatments is still in a condition that can be tolerated by tilapia, according to the statement Effendie (2003) which states that the optimal pH range for tilapia aquaculture is 6-8.5. Also in accordance with SNI 7550: 2009 the optimal pH for tilapia aquaculture is 6.5 - 8.5.

Temperature is a very important factor in fish farming, temperature can affect the body's metabolic rate in fish. The temperature range during the study for all treatments was around 25-29 °C according to the statement of Effendi et al. (2015) which states that the optimum temperature conditions for fish growth are 25-32 °C. This is also in accordance with SNI 7550: 2009 the optimal temperature for fish growth is 25-32 °C.

Dissolved oxygen is a very important factor in the cultivation of tilapia. Dissolved oxygen during the study from all treatments during the study 2.0 - 4.0 mg / l. Dissolved oxygen in all treatments during the study was still within the tolerance limits for tilapia, according to the statement (Popma and Masser, 1999) that tilapia can survive in the dissolved oxygen (DO) range of more than 0.3 mg / l. The value of dissolved oxygen for tilapia culture according to SNI 7550: 2009 states that the standard value of optimum levels of dissolved oxygen for the growth of tilapia is more than 3 mg / l.

Ammonia value in a fish culture container is very influential on the growth and survival of tilapia. Ammonia value at the beginning of the study was highest at P3 treatment and the lowest value at P0 treatment. Ammonia value at the end of the research was highest in the P0 treatment while the ammonia value of the other treatments was the same at 0.25 mg / l. The high value of ammonia at the beginning of the study was due to the high value of ammonia in tempe wastewater. The high value of ammonia is caused by the lack of ammonia content in the tempe waste by decomposing bacteria. As Harahap (2013) said, the ammonia in tempe wastewater will be broken down by bacteria (aerob-anaerobes) such as *Lactobacillus*, *Bifidobacterium*, *Clostridium*, *Bacteroides* and *Streptococcus* bacteria. According to Crab (2010).

CONCLUSIONS AND RECOMMENDATIONS

Conclusion

The conclusion that can be drawn from the research is the provision of tempeh liquid waste in the media of tilapia (*Oreochromis niloticus*) significantly influences ($P < 0.05$) on absolute growth but does not significantly affect the specific daily weight gain and absolute length as well as on feed efficiency.

Suggestion

In order to give maximum results, it is expected that there will be a researcher who is interested in continuing this research to find out whether or not the application of tempe liquid waste is true.

Acknowledgement

The author would like to thank Mr. Nurul as a tempe entrepreneur, who has provided tempe liquid waste as a treatment material in this study.

References

- Ahmadi, H., N. Iskandar and Kurniawati. 2012. Probiotics in Feeding the Growth of Sangkuriang Catfish (*Clarias* sp.) In Nursery II. Journal of Fisheries and Maritime Affairs. 3 (4): 99-107.
- Andersen JH, Schlüter L, Ærtebjerg G. 2006. Coastal eutrophication: recent developments in definitions and implications for monitoring strategies. Journal of Plankton Research. 28(7): 621-628.
- Anderson DM, Glibert PM, Burkholder JM. 2002. Harmful Algal Blooms and Eutrophication: Nutrient sources, composition, and Consequences. Estuaries. 25(4b): 704-726
- Agustin, Ruli, Ade DS, Yulisman. 2014. Konversi Pakan, Laju Pertumbuhan, Kelangsungan Hidup Dan Populasi Bakteri Benih Ikan Gabus (*Channa striata*) Yang Diberi Pakan Dengan Penambahan Probiotik. Jurnal Akuakultur Rawa Indonesia, 2 (1): 55- 66

Bodies Standart National. 2012. Tempe: From Indonesia to World. Jakarta. Pusido
Bodies Standart National

Crab, R., B. Chielens, M. Wille, P. Bossier and W. Verstraete. 2009. The Effect of
Different Carbon Sources on The Nutritional Value of Bioflocs, a Feed for
Macrobrachium rosenbergii Postlarvae. *Aquaculture Research*. 41: 559-
567

Dewantoro, GW 2001. Fecundity and larval production in betta fish
(*Bettasplendens* Regan), different age and natural food. Faculty
Biology, Jakarta National University. *Indonesian Iktiologi Journal*, 1. (2):
49-52.

Effendi, INJ Bugri, and Widanarni. 2006. The effect of stocking density on the
survival and growth of *Osphronemus gouramy* seeds *gouramy*. size 2 cm.
Indonesian Aquaculture Journal, 5 (2): 127-135.

Effendi, H. 2003. Water Quality Study. Kanisius. Yogyakarta

Gatesoupe, FJ 2008. Updating the Importance of Lactic Acid Bacteria in Fish
farming: natural occurrence and probiotic treatments. *J. Mol. Microbiol.*
Biotechnol., 14 (1-3): 107-114.

Effendi, H., BA Utomo, GM Darmawangsa, RE Karo-karo. 2015.
Phytoremediasilimbah cultivation of catfish (*Clarias* sp.) With kale
(*Ipomea aquatica*) and pakcoy (*Brassica rapa chinensis*) in the
recirculation system. *Ecolab*, 9 (2): 47-104.

Irianto, A., PAW Robertson and B. Austin, 2003. Oral administration of formalin
inactivated cells of *Aeromonas hydrophila* A3-51 controls infection by
atypical *A. salmonicida* in goldfish, *Carassius auratus* (L.). *Journal. of Fish*
Diseases, 26: 117-120.

Mashuri, Sumarjan, and Abidin, Z., 2012. The Effect Of Different Feed Types
On The Growth Of Eels (*Monopterus albus zuiewu*). *Research journal.*
Faculty University Agriculture Mataram

Ministry of Maritime Affairs and Fisheries. 2017. Marine and fisheries main
analysis and data by province in 2012. Central data, statistics and
information secretariat general of the ministry of fisheries and fisheries,
Jakarta

NRC. 1993. Nutrient Requirements of Warm Water Fishes and Shelfish. National
Academy of Sciences. Washington DC. 181 p.

Noviyanti Mz, Kadarwan Soewardi, NikenTunjung Murti Pratiwi. 2015.
Determination of Water Support Capacity for Natural Fisheries (Case Study:
Situ Cilala, Bogor Regency). *Indonesian Journal of Agricultural Sciences*
(JIPI). Vol. 20 (1): 66-71.

Said Idaman Nusa. 1999. Tofu-Tempe Wastewater Treatment Technology with Anaerobic and Aerobic Biofilter Processes. Jakarta: Directorate of Environmental Technology.

Sulistijowati S,R., Mile, L., 2016, Identification of lactic acid bacteria isolates from intestine of milkfish (*chanos-chanos*) potential activity against pathogen bacteria used pcr 18s rna methode, International jurnal bio-Science and Bio-Technology, 8(3), 127-134.

Wulandari, R., 2017, Kajian mikrobiologi penyakit ikan, UMRAH Press, Tanjungpinang

Popma, T., Masser, M. 1999. Tilapia life history and biology. Southern regionalaquaculture center publication no. 283.

Indonesian National Standard No. 7550: 2009. Production of Tilapia (*Oreochromis niloticus*) Enlargement Class in Calm Ponds. National Standardization Agency, Jakarta.

Sampe Harahap. 2013. Water Pollution Due to High Ammonia Levels From Tempe Liquid Waste. Journal of Aquatics Vol IV NO 2/2 September 2013 (183-194) ISSN 0853-2523

Widanarni, D., Wahjuningrum, and M. Setiawati. 2009. Optimization of Super-Intensive Culture of Environmentally Friendly Tilapia: Biofloc Microbial Dynamics. LPPM Seminar Material. Faculty of Fisheries and Marine Science. Bogor Agricultural Institute. 40 p.

Zakiyah Zahra Nur Amaliah, Saiful Bahri, and Princess Amelia. 2018. Isolation and Characterization of Lactic Acid Bacteria from Soy Bean Soaking Liquid Waste. JFFI. 2018; 5 (1) 253-257.

Lampiran



UNIVERSITAS MUHAMMADIYAH MALANG
FAKULTAS PERTANIAN-PETERNAKAN

Jurusan : Perikanan

Jl. Raya Tlogomas No. 246 Telp. (0341) 464318 psu 113 , 114 Malang – 65144
Fax. (0341) 460782 ; E-mail : fpp@umm.ac.id

FORM HASIL CEK PLAGIARISME LAPORAN TUGAS AKHIR

Nama Mahasiswa : Heryanto
NIM : 201610260311031
Judul TA : Pemanfaatan Limbah Cair Tempe dalam Media Air Budidaya untuk
Pertumbuhan dan Efisiensi Pakan Ikan Nila (*Oreochromis niloticus*)

Hasil Cek Plagiarisme dengan Turnitin

No	Komponen Pengecekan	Maksimum Kesamaan (%)	Hasil Cek plagiarisme (%)
1	Naskah publikasi	25	0

Malang, 27 Agustus 2020

Mengetahui
Ketua Program Studi Akuakultur

Ganjar Adhywirawan Sutarjo, S.Pi, M.P

Admin Turnitin
Program Studi Akuakultur

Anis Zubaidah, S.Pi, M.Si